

Azathioprine factsheet

Azathioprine is a drug that was originally developed to prevent graft rejection in transplant patients. It has been available since the 1960s and has been used for many years to treat severe eczema. Clinical trials in the 1990s provided good evidence that azathioprine is effective for resistant moderate-to-severe disease. It has also frequently been used in other skin diseases as a means of reducing oral steroid use. However, azathioprine is not licensed for use in atopic eczema and is prescribed 'off license' only by a specialist doctor such as a dermatologist.

How does azathioprine work?

Azathioprine is an immunosuppressant drug that is also known as an antimetabolite. It interferes with the growth of certain types of white blood cells (lymphocytes) that are involved in creating the inflammation associated with eczema. It takes a little longer to work compared to ciclosporin and benefits are usually not seen until after 4-6 weeks after starting treatment, so it is less suitable for acute flares because it takes longer to take effect. Further improvements then occur over the next few months.

When is azathioprine used?

Azathioprine is mainly used in the UK to treat people with severe atopic eczema that is unresponsive to normal topical treatment. The reasons for prescribing azathioprine are similar to those for ciclosporin, although it is less suitable for acute flares because it takes longer to take effect. Azathioprine is available as 25 mg and 50 mg tablets. For patients with normal thiopurine methyltransferase (see opposite), azathioprine can be taken as a single dose. But if side effects such as sickness occur, dividing the dose and taking twice a day may help. A starting dose of 2 mg/kg is often used initially, and then the dose is gradually increased according to the response. The usual maintenance dose is between 100 mg and 250 mg per day (2.5 mg/kg per day).

What are the side effects?

It is important to remember that most patients do not experience significant side effects. Blood tests will be taken before treatment to assess the individual risk for developing side effects. The main side effect concern with azathioprine is bone-marrow suppression. This can result in severe anaemia and risk of infection. A blood test is now available to determine which patients are most at risk of developing this side effect with azathioprine. About one in 200 people have very low levels of an enzyme called thiopurine methyltransferase (TPMT). They are unable to break down azathioprine in the normal way and are at high risk of dangerous bone-marrow suppression. **These patients should not receive azathioprine.** A further group of patients who have intermediate activity of TPMT may be given azathioprine, but it is recommended that a reduced dose be given, starting at 0.5 mg/kg and increasing to 1.0 mg/kg. If the level of TPMT is normal, risk of developing side effects is low. All patients taking azathioprine require regular blood tests before and during treatment, to check for any signs of bone-marrow suppression even if their TPMT level is normal. You should seek medical advice if you become unwell or develop signs of infection. It is important to report any unusual bruising or bleeding, which may be a sign that the bone marrow is being affected.

Azathioprine factsheet

Azathioprine can sometimes cause problems with the liver. Liver function tests are therefore performed at the start of treatment, with weekly monitoring to begin with, then every 3 months if the test results are stable. There is also a theoretical concern that long-term treatment may increase the risk of certain types of malignancy but there is no evidence that this is the case in the short-to-medium term. Nausea, diarrhoea and loss of appetite may be a problem for some patients, but this usually responds to a reduction in dose. Hair loss can occur, but this is usually mild and reversible on stopping the drug. Occasionally a drug hypersensitivity syndrome may develop – this feels like ‘flu’ with aches, pains and fever. (below). If this occurs the drug should be stopped immediately and the prescribing doctor informed.

Can azathioprine be used in combination with other drugs?

Certain tablets increase the risk of side effects with azathioprine and should therefore be avoided. If any new tablets are considered, it is important to let the doctor know you are taking azathioprine. In particular, allopurinol – a drug that is used to treat gout – can cause severe bone-marrow suppression. Warfarin (used to thin the blood), and some antibiotics (such as Septrin and trimethoprim) may have a similar effect and should be avoided. Some blood pressure tablets (those that are ACE inhibitors) should also be avoided.

What blood tests are needed?

All patients taking azathioprine require blood tests before and during treatment to check for any signs of bone-marrow suppression, even if their TPMT level is normal. As azathioprine can sometimes cause problems with the liver, blood count and liver function tests will be carried out weekly until a maintenance dose has been established, then blood tests will take place every 3 months.

Who is prescribed azathioprine?

Azathioprine can be prescribed for adults and older children with severe eczema, but only under consultant dermatologist supervision.

Both men and women must stop taking azathioprine for at least 3 months before trying to conceive. The effectiveness of intrauterine contraceptive devices may be affected by azathioprine, so other or additional forms of contraception are recommended.

Azathioprine should not be used in pregnancy without careful assessment of the risks and benefits.

Those taking azathioprine are advised not to breastfeed.

Immunisations and azathioprine

‘Live’ vaccines are not recommended whilst taking azathioprine. These include:

- *Oral polio drops.* Polio injections are inactive and safe; other members of your household should also be given inactive, rather than live, polio vaccine.
- *Nasal ‘flu’ (influenza) vaccine.* Yearly influenza and 5-yearly pneumococcal injections are safe and should ideally be given to everybody on azathioprine.
- *MMR (mumps, measles and rubella).*
- *Oral typhoid.*
- *BCG for tuberculosis (TB).*
- *Chickenpox (Varicella zoster).* If you have never had chickenpox, your doctor may recommend vaccination against this before starting azathioprine. If this was not possible and you come into contact with a person with chickenpox or shingles whilst you are taking azathioprine, you should go to your doctor straightaway as you may need special preventative treatment.

continued on page 3

Azathioprine factsheet

- *Shingles (Herpes zoster).*
- *Yellow fever.*

If you have any questions or concerns, please do not hesitate to speak to a doctor, pharmacist or nurse caring for you.

DISCLAIMER

Our publications contain information and general advice about eczema. They are written and reviewed by dermatology experts, with input from people with eczema. We hope you find the information helpful, although it should not be relied upon as a substitute for personalised advice from a qualified healthcare professional. While we strive to ensure the information is accurate and up-to-date, National Eczema Society does not accept any liability arising from its use. We welcome reader feedback on our publications, please email us at info@eczema.org

Factsheet last reviewed October 2018.

© National Eczema Society, June 2019. All rights reserved. Except for personal use, no part of this work may be distributed, reproduced, downloaded, transmitted or stored in any form without the written permission of National Eczema Society.



National Eczema Society is the UK charity for everyone affected by eczema. We help support people with eczema, providing information and advice, which we deliver through our website, social media, campaigns, publications and nurse-supported Helpline. We also provide a voice for people with eczema, raising awareness of the condition, supporting research into new treatments and campaigning for better medical care.

National Eczema Society is a registered charity in England and Wales (No. 1009671) and in Scotland (No. SC043669). Registered Office: 11 Murray Street, London NW1 9RE

National Eczema Society Helpline: 0800 089 1122 Email: helpline@eczema.org www.eczema.org